

Local Coverage Determination (LCD): MoIDX: NRAS Genetic Testing (L36335)

Links in PDF documents are not guaranteed to work. To follow a web link, please use the MCD Website.

Contractor Information

Contractor Name	Contract Type	Contract Number	Jurisdiction	State(s)
Noridian Healthcare Solutions, LLC	A and B MAC	01111 - MAC A	J - E	California - Entire State
Noridian Healthcare Solutions, LLC	A and B MAC	01112 - MAC B	J - E	California - Northern
Noridian Healthcare Solutions, LLC	A and B MAC	01182 - MAC B	J - E	California - Southern American Samoa
Noridian Healthcare Solutions, LLC	A and B MAC	01211 - MAC A	J - E	Guam Hawaii Northern Mariana Islands American Samoa
Noridian Healthcare Solutions, LLC	A and B MAC	01212 - MAC B	J - E	Guam Hawaii Northern Mariana Islands
Noridian Healthcare Solutions, LLC	A and B MAC	01311 - MAC A	J - E	Nevada
Noridian Healthcare Solutions, LLC	A and B MAC	01312 - MAC B	J - E	Nevada American Samoa California - Entire State
Noridian Healthcare Solutions, LLC	A and B MAC	01911 - MAC A	J - E	Guam Hawaii Nevada Northern Mariana Islands

[Back to Top](#)

LCD Information

Document Information

LCD ID L36335	Original Effective Date For services performed on or after 07/05/2016
LCD Title MoIDX: NRAS Genetic Testing	Revision Effective Date For services performed on or after 01/19/2017
AMA CPT / ADA CDT / AHA NUBC Copyright Statement CPT only copyright 2002-2017 American Medical Association. All Rights Reserved. CPT is a registered trademark of the American Medical Association. Applicable FARS/DFARS Apply to Government Use. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.	Revision Ending Date N/A
	Retirement Date N/A
	Notice Period Start Date 05/19/2016
	Notice Period End Date 07/04/2016

The Code on Dental Procedures and Nomenclature (Code) is published in Current Dental Terminology (CDT). Copyright © American Dental Association. All rights reserved. CDT and CDT-2016 are trademarks of the American Dental Association.

UB-04 Manual. OFFICIAL UB-04 DATA SPECIFICATIONS MANUAL, 2014, is copyrighted by American Hospital Association ("AHA"), Chicago, Illinois. No portion of OFFICIAL UB-04 MANUAL may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior express, written consent of AHA." Health Forum reserves the right to change the copyright notice from time to time upon written notice to Company.

CMS National Coverage Policy Title XVIII of the Social Security Act, §1862(a)(1)(A). Allows coverage and payment for only those services that are considered to be reasonable and necessary.

Title XVIII of the Social Security Act, §1862(a)(1)(D). Allows coverage and payment for clinical care items and services provided with the concurrence of the Secretary and with respect to research and experimentation conducted by, or under contract with, the Medicare Payment Advisory Commission or the Secretary, which are not reasonable and necessary to carry out the purposes of section

Title XVIII of the Social Security Act, §1833(e). Prohibits Medicare payment for any claim which lacks the necessary information to process the claim.

42 CFR 410.32(a). States diagnostic tests must be ordered by the physician treating the beneficiary.

CMS On-Line Manual, Publication 100-02, Medicare Benefit Policy Manual, Chapter 15, §§ 80.6, Requirements for Ordering and Following Orders for Diagnostic Tests.

CMS On-Line Manual, Publication 100-08, Medicare Benefit Policy Manual, Chapter 3, §§ 3.4.1.3, Diagnosis Code Requirements.

Coverage Guidance

Coverage Indications, Limitations, and/or Medical Necessity

Indications:

This is limited coverage policy for genetic testing of tumor tissue for somatic mutations in the NRAS gene (81311). Noridian will cover NRAS testing for metastatic colorectal cancer, per NCCN guidelines (Version 2.2016).

All other NRAS testing is non-covered.

Background:

RAS oncogene is a superfamily of signal transduction proteins, which are proteins that communicate signals between the cells. DNA mutations in the RAS family genes turns the signals on permanently such that the cells divide nonstop, leading to cancer. Three of this family's proteins, HRAS, KRAS, and NRAS are important in tumors and encode 21kD proteins called p21s.

Previous studies have shown that targeting oncogenic NRAS-driven melanomas requires decrease in both pERK and pAKT downstream of RAS-effectors for efficacy, which could be achieved by either targeting both BRAF and CRAF or BRAF and PIK3CA simultaneously in NRAS mutant tumor cells.

Colorectal Cancer:

Multiple signaling pathways are involved in colorectal cancer pathogenesis. The epidermal growth factor receptor (EGFR) plays a key role in activation of these pathways and is commonly overexpressed in metastatic colorectal cancer (mCRC). Consequently, EGFR is a target of anticancer therapies. Two of these drugs, cetuximab and panitumumab, are monoclonal antibodies that block EGFR action. The 2013 NCCN Clinical Practice Guidelines for Colon Cancer describes a recent study by Douillard et al [2013] which reported that 17% of 641 patients from the

PRIME trial without KRAS exon 2 mutations were found to have mutations in exons 3 and 4 of KRAS or mutations in exons 2, 3, and 4 of NRAS. A predefined retrospective analysis of a subset of these patients showed that progression free survival (PFS) and overall survival (OS) were decreased in those who received panitumumab plus FOLFOX compared to those who received FOLFOX alone. For this reason the FDA indication for panitumumab was recently updated to state that panitumumab is not indicated for the treatment of patients with NRAS mutation-positive disease in combination with oxaliplatin-based chemotherapy.

In chemotherapy-refractory patients, fewer than 10% of patients who harbor one of these mutations respond to EGFR immunotherapy. The American Society of Clinical Oncology (ASCO) and the National Comprehensive Cancer Network (NCCN) both recommend KRAS mutation testing prior to prescribing EGFR antagonist therapy for patients with mCRC and state that alternative therapy should be prescribed when mutations are detected.

However, NCCN Colorectal Guidelines (Version 2.2016) recommend "All patients with metastatic colorectal cancer should have tumor tissue genotyped for RAS mutations (KRAS and NRAS) and BRAF mutations. Patients with any known KRAS mutation (exon 2 or non- exon 2) or NRAS mutation should not be treated with either cetuximab or panitumumab." Evidence increasingly suggests that BRAF V600E mutation makes response to panitumumab or cetuximab highly unlikely, as a single agent, or in combination with cytotoxic chemotherapy. In light of the above, KRAS, NRAS and BRAF are covered for metastatic colorectal cancer.

Metastatic Melanoma:

The NRAS gene encodes a protein that helps control cell division. Approximately 15% to 20% of melanomas harbor an oncogenic NRAS mutation. NRAS mutations can occur in all melanoma subtypes, but may be slightly more common in skin with chronic sun damage or in nodular melanomas. In addition, NRAS mutations are not found in tumors with BRAF mutations.

Several studies have been carried out to examine whether mutations in BRAF and NRAS confer different pathological features and clinical behavior. The effect of these mutations on clinical outcome remains uncertain with previous studies reporting conflicting results.

(Per NCCN Guidelines 3.2016- BRAF- targeted Therapies: "Approximately half of patients with metastatic cutaneous melanoma harbor an activating mutation of BRAF, an intracellular signaling kinase in the MAPK pathway. Most BRAF-activating mutations occurring in melanomas are at residue V600, usually V600E but occasionally V600K or other substitutions. BRAF inhibitors have been shown to have clinical activity in melanomas with BRAF V600 mutations. Inhibitors of MEK, a signaling molecule downstream of BRAF, may potentiate these effects. Recent efficacy and safety data from large randomized trials testing BRAF and MEK inhibitors have significantly impacted the recommended treatment options for patients with BRAF-mutation positive advanced melanoma.")

The NRAS protein is a GTPase which can lead to the activation of other proteins (such as AKT and MEK) that are also in pathways that help regulate cell division. In theory, drugs that inhibit AKT or MEK also have the potential to counteract the effects of NRAS mutations, although NRAS targeting therapies are still in clinical trials. In addition, pathways that help regulate cell division also include other proteins that could potentially be targeted such as PI3K and mTOR.

Melanomas can be tested for NRAS mutations with targeted sequencing. There are several manufacturers of targeted genetic tests that can detect NRAS mutations in melanoma tumor samples. The prognostic significance of NRAS mutations is still not well understood and further investigation of the histologic types of melanoma with specific NRAS mutations in a larger series is necessary to validate these apparent impacts on patient outcomes. In smaller subsets of cutaneous melanoma, other activating mutations have been described, including NRAS, c-KIT, and CDK4.

Other Cancers:

Other neoplastic diseases in which NRAS mutations have been reported in the primary literature include: myeloid leukemia, bladder cancer, liver cancer, and proliferative thyroid lesions.

Schulten et al [2-013] directly sequenced mutational hotspot regions encompassing codons 12, 13, and 61 of the RAS genes in 381 cases of thyroid lesions. In addition, the putative NRAS hotspot region encompassing codon 97 was sequenced in 36 thyroid lesions. Schulten and team found mutations in 16 out of 57 patients.

Kompier et al [2010] reports that although they have been reported, NRAS mutations are not common in bladder cancer.

Although NRAS mutations have been identified in the above tumor types, evidence in the primary literature is limited with regard to the clinical utility of NRAS mutation testing and its impact on management and survival. There is currently insufficient evidence to demonstrate clinical utility of NRAS testing in these tumor types.

NRAS Testing in relation to Noonan syndrome diagnosis:

Noonan syndrome is a common autosomal dominant condition with an incidence of 1/1,000 to 1/2,500 people. Unlike the somatic tumor mutations discussed above, Noonan syndrome may be caused by a germline mutation in the NRAS gene which would be present in every cell of the body. Noonan syndrome is characterized by a number of phenotypic findings including distinctive facial features, short stature, heart defects, cryptorchidism, lymphedema, and coagulation defects, among others. Several syndromes have features that overlap clinically with Noonan syndrome including cardiofaciocutaneous syndrome, Costello syndrome, LEOPARD syndrome and Noonan--like syndrome with loose anagen hair. The genetic etiologies of these conditions can also overlap with Noonan syndrome.

Several of these disorders have been referred to as neurocardiofacialcutaneous syndromes, RASopathies or Ras/MAPK pathway disorders and have a shared pathway of genetic function.

They are characterized by facial dysmorphism, cardiac disease, reduced growth, skeletal and ectodermal defects and variable cognitive deficits. They also share a predisposition to development of malignancies.

Overall, approximately 75% of individuals with Noonan syndrome will have an identifiable mutation with gene panel testing. To date, NRAS mutations have been found in four individual case reports which suggests that NRAS testing for Noonan syndrome is unlikely to yield positive results. The clinical features appear to be typical with no particular or distinctive phenotype observed suggesting that mutation testing targeted to select individuals is not feasible.

Genotype-/phenotype correlations have emerged that can help to direct medical management for those affected with an associated condition, but not specifically for NRAS mutations. For instance, mutations in the SOS1 gene have been associated with an increased chance for ectodermal involvement, development of certain solid tumors, pulmonary stenosis, and atrial and ventricular septal defects; with an associated decreased prevalence of cognitive defects, short stature, and hypertrophic cardiomyopathy.

Medical management recommendations are available for many of the Noonan syndrome spectrum disorders. Overlapping features result in overlapping medical management recommendations, typically guided by clinical features.

[Back to Top](#)

Coding Information

Bill Type Codes:

Contractors may specify Bill Types to help providers identify those Bill Types typically used to report this service. Absence of a Bill Type does not guarantee that the policy does not apply to that Bill Type. Complete absence of all Bill Types indicates that coverage is not influenced by Bill Type and the policy should be assumed to apply equally to all claims.

N/A

Revenue Codes:

Contractors may specify Revenue Codes to help providers identify those Revenue Codes typically used to report this service. In most instances Revenue Codes are purely advisory. Unless specified in the policy, services reported under other Revenue Codes are equally subject to this coverage determination. Complete absence of all Revenue Codes indicates that coverage is not influenced by Revenue Code and the policy should be assumed to apply equally to all Revenue Codes.

N/A

CPT/HCPCS Codes

Group 1 Paragraph: N/A

Group 1 Codes:

81311 NRAS (NEUROBLASTOMA RAS VIRAL [V-RAS] ONCOGENE HOMOLOG) (EG, COLORECTAL CARCINOMA), GENE ANALYSIS, VARIANTS IN EXON 2 (EG, CODONS 12 AND 13) AND EXON 3 (EG, CODON 61)
81479 UNLISTED MOLECULAR PATHOLOGY PROCEDURE

ICD-10 Codes that Support Medical Necessity

Group 1 Paragraph: Although not specifically addressed in the ICD-10-CM Official Guidelines for Coding and Reporting 2016, when an encounter is for management of a complication associated with a neoplasm (NRAS testing for metastatic colon cancer), the complication (metastasis) is coded first, followed by the appropriate codes for the neoplasm.

*Primary Diagnoses are listed in Group 1 and Secondary Diagnoses in Group 2.

Group 1 Codes:

ICD-10 Codes

Description

C77.0	Secondary and unspecified malignant neoplasm of lymph nodes of head, face and neck
C77.1	Secondary and unspecified malignant neoplasm of intrathoracic lymph nodes
C77.2	Secondary and unspecified malignant neoplasm of intra-abdominal lymph nodes
C77.3	Secondary and unspecified malignant neoplasm of axilla and upper limb lymph nodes
C77.4	Secondary and unspecified malignant neoplasm of inguinal and lower limb lymph nodes
C77.5	Secondary and unspecified malignant neoplasm of intrapelvic lymph nodes
C77.8	Secondary and unspecified malignant neoplasm of lymph nodes of multiple regions
C77.9	Secondary and unspecified malignant neoplasm of lymph node, unspecified
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.1	Secondary malignant neoplasm of mediastinum
C78.2	Secondary malignant neoplasm of pleura
C78.30	Secondary malignant neoplasm of unspecified respiratory organ
C78.39	Secondary malignant neoplasm of other respiratory organs
C78.4	Secondary malignant neoplasm of small intestine
C78.5	Secondary malignant neoplasm of large intestine and rectum
C78.6	Secondary malignant neoplasm of retroperitoneum and peritoneum
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C78.80	Secondary malignant neoplasm of unspecified digestive organ
C78.89	Secondary malignant neoplasm of other digestive organs
C79.00	Secondary malignant neoplasm of unspecified kidney and renal pelvis
C79.01	Secondary malignant neoplasm of right kidney and renal pelvis
C79.02	Secondary malignant neoplasm of left kidney and renal pelvis
C79.10	Secondary malignant neoplasm of unspecified urinary organs
C79.11	Secondary malignant neoplasm of bladder
C79.19	Secondary malignant neoplasm of other urinary organs
C79.2	Secondary malignant neoplasm of skin
C79.31	Secondary malignant neoplasm of brain
C79.32	Secondary malignant neoplasm of cerebral meninges
C79.40	Secondary malignant neoplasm of unspecified part of nervous system
C79.49	Secondary malignant neoplasm of other parts of nervous system
C79.51	Secondary malignant neoplasm of bone
C79.52	Secondary malignant neoplasm of bone marrow
C79.60	Secondary malignant neoplasm of unspecified ovary
C79.61	Secondary malignant neoplasm of right ovary
C79.62	Secondary malignant neoplasm of left ovary
C79.70	Secondary malignant neoplasm of unspecified adrenal gland
C79.71	Secondary malignant neoplasm of right adrenal gland
C79.72	Secondary malignant neoplasm of left adrenal gland
C79.81	Secondary malignant neoplasm of breast
C79.82	Secondary malignant neoplasm of genital organs
C79.89	Secondary malignant neoplasm of other specified sites
C79.9	Secondary malignant neoplasm of unspecified site

Group 2 Paragraph: Secondary Diagnoses:

Group 2 Codes:

ICD-10 Codes	Description
C18.0	Malignant neoplasm of cecum
C18.1	Malignant neoplasm of appendix
C18.2	Malignant neoplasm of ascending colon
C18.3	Malignant neoplasm of hepatic flexure
C18.4	Malignant neoplasm of transverse colon
C18.5	Malignant neoplasm of splenic flexure
C18.6	Malignant neoplasm of descending colon
C18.7	Malignant neoplasm of sigmoid colon
C18.8	Malignant neoplasm of overlapping sites of colon
C18.9	Malignant neoplasm of colon, unspecified
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum

ICD-10 Codes that DO NOT Support Medical Necessity N/A

ICD-10 Additional Information

[Back to Top](#)

General Information

Associated Information

N/A

Sources of Information and Basis for Decision

1. Akslen LA, Angelini S, Straume O, et al. BRAF and NRAS mutations are frequent in nodular melanoma but are not associated with tumor cell proliferation or patient survival. *J Invest Dermatol*. 2005 Aug;125(2):312-7.
2. Allegra CJ, Jessup JM, Somerfield MR, et al. American Society of Clinical Oncology provisional clinical opinion: testing for KRAS gene mutations in patients with metastatic colorectal carcinoma to predict response to anti-epidermal growth factor receptor monoclonal antibody therapy. *J Clin Oncol*. 2009;27:2091--2096.
3. Benson AB, et al. NCCN Clinical Practice Guidelines in Oncology™: colon cancer. National Comprehensive Cancer Network Web site. http://www.nccn.org/professionals/physician_gls/PDF/colon.pdf . Version 2. 2014. Accessed November 22, 2013.
4. Cirstea IC, Kutsche K, Dvorsky R, et al. A restricted spectrum of NRAS mutations causes Noonan syndrome. *Nat Genet*. 2010;42:27-9.
5. DeRoock W, Claes B, Bernasconi D, et al. Effects of KRAS, BRAF, NRAS, and PIK3CA mutations on the efficacy of cetuximab plus chemotherapy in chemotherapy-refractory metastatic colorectal cancer: A retrospective consortium analysis. *Lancet Oncol*. 2010;11:753--762.
6. Devitt B, Liu W, Salemi R, et al. Clinical outcome and pathological features associated with NRAS mutation in cutaneous melanoma. *Pigment Cell Melanoma Res*. 2011; 24: 666-672.
7. Douillard JY, Oliner KS, Siena S, et al. Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer. *N Engl J Med* 2013;369:1023--1034.
8. Edlundh--Rose E, Egyházi S, Omholt K, et al. NRAS and BRAF mutations in melanoma tumours in relation to clinical characteristics: a study based on mutation screening by pyrosequencing. *Melanoma Res* 16 (6): 471--8, 2006.
9. Gavin PG, Colangelo LH, Fumagalli D, et al. Mutation profiling and microsatellite instability in stage II and III colon cancer: an assessment of their prognostic and oxaliplatin predictive value. *Clin Cancer Res*. 2012 Dec 1;18(23):6531-41.
10. Goel VK, Lazar AJ, Warneke CL, et al. Examination of mutations in BRAF, NRAS, and PTEN in primary cutaneous melanoma. *J Invest Dermatol* 126 (1): 154--60, 2006.

11. Jaiswal BS, Janakiraman V, Kljavin NM, et al. Combined Targeting of BRAF and CRAF or BRAF and PI3K Effector Pathways Is Required for Efficacy in NRAS Mutant Tumors. *Plos one* 2009;4(5) e5717.
12. Kakavand H, Scolyer RA, Thompson JF, Mann GJ. Identification of new prognostic biomarkers for Stage III metastatic melanoma patients. *OncoImmunology* 2013;2(9).
13. Kompier LC, Lurkin I, Van der Aa MN, van Rhijn BWG, van der Kwast, Zwarthoff EC. FGFR3, HRAS, KRAS, NRAS and PIK3CA Mutations in Bladder Cancer and Their Potential as Biomarkers for Surveillance and Therapy. *Plos One* 2010;5(11).
14. Lee JH, Choi JW, Kim YS. Frequencies of BRAF and NRAS mutations are different in histological types and sites of origin of cutaneous melanoma: a meta- analysis. *British Association of Dermatologists*. 2011;164: 776–784.
15. Lepri F, De Luca A, Stella L, et al. SOS1 mutations in Noonan syndrome: molecular spectrum, structural insights on pathogenic effects, and genotype/phenotype correlations. *Hum Mutat*. 2011 Jul;32(7):760--72.
16. NCCN Guidelines. Version 3.2014.
https://www.nccn.org/store/login/login.aspx?ReturnURL=http://www.nccn.org/professionals/physician_gls/pdf/melanoma.pdf
17. Schulten HJ, Salama S, Al-Ahmadi A, et al. Comprehensive Survey of HRAS, KRAS, and NRAS Mutations in Proliferative Thyroid Lesions from An Ethnically Diverse Population. *Anticancer Res*. 2013 Nov;33(11):4779-84.
18. Singhal R and Kandel ES. The response to PAK1 inhibitor IPA3 distinguishes between cancer cells with mutations in BRAF and Ras oncogenes. *Oncotarget* 2012; 3(7):700-708.
19. Stockman D, Tetzlaff MT, AlZaid T, et al. Differential clinical associations of BRAF and NRAS mutations among histologic types of cutaneous melanomas. *J Clin Oncol* 2013 (suppl; abstr e20034).
20. Tartaglia M, Gelb BD. Disorders of dysregulated signal traffic through the RAS--MAPK pathway: phenotypic spectrum and molecular mechanisms. *Ann N Y Acad Sci*. 2010 Dec;1214:99--121.

21. Tartaglia M, Gelb BD, Zenker M. Noonan syndrome and clinically related disorders. *Best Pract Res Clin Endocrinol Metab.* 2011 Feb;25(1):161-79.
22. Tidyman WE, Rauen KA. The RASopathies: developmental syndromes of Ras/MAPK pathway dysregulation. *Curr Opin Genet Dev.* 2009 Jun;19(3):230-6.
23. Vaughn CP, ZoBell SD, Furtado LV, Baker CL, Samowitz WS. Frequency of KRAS, BRAF, and NRAS mutations in colorectal cancer. *Genes, Chromosomes and Cancer*, 2011;50(5):307--312.

Revision History Information

Revision History Date	Revision History Number	Revision History Explanation	Reason(s) for Change
01/19/2017	R2	Added the following ICD-10 codes under Group 1 that were left out of the C79.00-C79.9 range in error. C79.01, C79.02, C79.10,C79.11, C79.19, C79.2, C79.31, C79.32, C79.40, C79.49, C79.51, C79.52, C79.60, C79.61, C79.62, C79.70, C79.71, C79.72, C79.81, C79.82, C79.89, C79.9 Added primary and secondary diagnoses codes under ICD-10 Codes that Support Medical Necessity:	<ul style="list-style-type: none">Revisions Due To ICD-10-CM Code Changes
01/19/2017	R1	Group 1: C77.0-C77.9 Secondary and unspecified malignant neoplasm of lymph nodes C78.00-C78.89 – Secondary malignant neoplasm of respiratory and digestive organs C79.00-C79.9 Secondary malignant of other and unspecified sites Group 2: C18.0-C18.9 – Malignant neoplasm of colon C19 – Malignant neoplasm of recto-sigmoid junction C20 – Malignant neoplasm of rectum	<ul style="list-style-type: none">Revisions Due To ICD-10-CM Code Changes

Associated Documents

Attachments N/A

Related Local Coverage Documents Article(s) [A55048 - Response to Comments: MoIDX: NRAS Genetic Testing](#)
LCD(s) [DL36308](#) - (MCD Archive Site) [DL36335](#) - (MCD Archive Site)

Related National Coverage Documents N/A

Public Version(s) Updated on 01/26/2017 with effective dates 01/19/2017 - N/A [Updated on 01/23/2017 with effective dates 01/19/2017 - N/A](#) [Updated on 05/03/2016 with effective dates 07/05/2016 - N/A](#) [Back to Top](#)

Keywords

- NRAS
- KRAS
- HRAS
- Noonan
- 81311
- 81479
- neoplasm
- melanoma
- MoIdx
-
-
-
-

Read the [LCD Disclaimer](#) [Back to Top](#)